


The control of algal abundance on coral reefs through the reintroduction of *Diadema*
antillarum



Stacey M. Williams, Ph.D.
Institute for Socio-Ecological Research
PO Box 3151
Lajas, PR 00667

December 5, 2018

(Cooperative Agreement NA17NOS4820037)



Executive Summary

In Puerto Rico, a more recent and potential threat to the coral reefs is the increasing abundance of a red encrusting algae, *Ramicrusta* sp. *Ramicrusta* is an encrusting red alga from the Peyssonneliaceae family (Rhodophyta). *Ramicrusta* was observed at 70% of the monitoring sites (42 sites) in the 2017-18 DNER's Puerto Rico Coral Reef Monitoring Program (PRCRMP) surveys. Moreover, *Ramicrusta* sp. was the main benthic substrate observed on shallow-water coral reefs at a majority of the reefs surveyed in Culebra Island and Fajardo. *Diadema antillarum*, a keystone herbivore, has been witnessed eating *Ramicrusta* (Williams pers comm, Ruiz 2015, Williams 2017) and in the field we observed a noticeably lower cover of *Ramicrusta* in areas where *D. antillarum* were present. The goal of this project was to increase herbivory rates on two coral reefs in Fajardo and decrease algal abundance, specifically *Ramicrusta* sp. In addition, it is also expected, that effective recruitment rates will increase as an indirect consequence of *D. antillarum* herbivory effect.

A total of 3,179 *D. antillarum* settlers were collected during the summer of 2017. They were brought back to the laboratory at the University of Puerto Rico and grown to a young adult size (test diameter between 3cm to 4cm). The survivorship of the individuals grown in laboratory was impacted by the passing of Hurricane Irma and Maria. *D. antillarum* settlers had to be placed in cages under the dock at the university because there was no electricity for three months. On January 4, 2018, 505 sea urchins were removed from the cages and placed back in the raceways outside (37% survivorship).

On August 22, 2018, at 1:00 PM, 480 young adults (3-4cm) were transferred to Cayo Diablo (240 in total) and Los Lobos (240 in total). Twenty-five urchins were placed in six corrals that were installed at each reef. Tops were installed on three of the corrals to fully enclose (hereafter closed) the area, and three corrals did not have tops. Most of the *D. antillarum* escaped the corrals a day after the restocking. Overall, the retention of individuals in corrals was greater in the closed corrals than in the open corrals. After the first month, an average of 44% and 52% of the restocked individuals were recorded in fully closed corrals in Los Lobos and Cayo Diablo, respectively. The majority of the corrals were completely destroyed and open by the second month, therefore the sea urchins were no longer confined. Therefore, the numbers of *D. antillarum* located in the corrals decreased even more so by the second month of monitoring.

The benthic composition did not vary at the control stations through the monitoring period. Therefore, it can be assumed that the changes observed in the benthic cover inside the corrals, both open and closed, are due to the grazing of restocked *D. antillarum*. One week after the restocking, the benthic substrate was characterized by more turf algae, and less *Ramicrusta*. There was a shift in benthic community structure after one and two months after the restocking (9/22/2018, 10/23/2018). The difference in benthic assemblage after one month was due to the reduction of *Ramicrusta* cover. The abundance of *Ramicrusta* was significantly reduced over the monitoring period (Indices 3-4). The reduction in *Ramicrusta* was slightly more evident at Cayo Diablo than at Los Lobos; where *Ramicrusta* was reduced as much as 63% at Cayo Diablo (Fig. 16) and 61% at Los Lobos (See Indices 5 and 6). After one and two months, the substrate inside the corrals was characterized by more available space and turf algae, which allows the settlement and colonization of corals and other benthic organisms. The evidence presented in this study supports that the action

of restocking of *Diadema antillarum* may be useful mitigation tool to control *Ramicrusta* and other algal abundance on the coral reefs.

Contents

Executive Summary	2
Figure List	6
Table List	7
Index List	8
Introduction.....	9
Methods.....	12
Collection	12
Grow-out	13
Restocking.....	14
Statistics	16
Results and Discussion	17
Collection	17
Grow-out	18
Restocking.....	18
Survivorship.....	18
Benthic Cover	22
Conclusions and recommendations.....	32
Acknowledgements.....	33
Indices	34
Literature review	43

Figure List

Figure 1 .Settlement sites in La Parguera, Puerto Rico	12
Figure 2 Astroturf or artificial turf used as settlement.....	12
Figure 3 Diadema antillarum settler	13
Figure 4 Wet laboratory at the University of Puerto Rico, Mayaguez.	13
Figure 5 Map of the restoration sites, Cayo Diablo and Los Lobos in Fajardo, Puerto Rico.	15
Figure 6 Pictures of the different corrals in Cayo Diablo (top six pictures) and in Los Lobos (bottom six), in Fajardo, Puerto Rico. Corral number is arranged from left to right.....	16
Figure 7 Pictures of the lab-reared Diadema antillarum restocked in the corral at Cayo Diablo and Los Lobos reefs in Fajardo, Puerto Rico.....	19
Figure 8 Photographs taken during the two-month monitoring showing the destructions of corrals at Cayo Diablo and Los Lobos reefs in Fajardo, Puerto Rico.	22
Figure 9 Photographs at the control station in Cayo Diablo (two photographs above random quadrat) and in Los Lobos (two photographs below fixed quadrats) in Fajardo, Puerto Rico.	23
Figure 10 The mean cover of benthic organisms in the corral type (open and closed) and quadrat type (fixed and random) through the sampling events at Cayo Diablo, Fajardo.	24
Figure 11 The mean cover of benthic organisms in the corral type (open and closed) and quadrat type (fixed and random) through the sampling events at Los Lobos, Fajardo.	25
Figure 12 Photographs of the fixed quadrats before and one week after the restocking of Diadema antillarum at Cayo Diablo (first two sets of photographs) and Los Lobos (third set of photograph) in Fajardo, Puerto Rico.....	28
Figure 13 Photographs taken to highlight the grazing of restored Diadma antillarum one week after the restocking at Cayo Diablo and Los Lobos in Fajardo, Puerto Rico.	29
Figure 14 Principal Coordinate Analysis on the benthic composition of closed (left) and open (right) corrals at Cayo Diablo during the sampling periods. Algae represents all macroalgae. ...	30
Figure 15 Principal Coordinate Analysis on the benthic composition of closed (left) and open (right) corrals at Los Lobos during the sampling periods. Algae represents all macroalgae.....	30
Figure 16 Photographs of a fixed quadrat in Corral 5 at Cayo Diablo before, one week and two months after the restocking of Diadema antillarum.....	31

Table List

Table 1. The total amount of settlers (#) collected at each sites from June to September 2017...	18
Table 2 The number of restocked <i>Diadema antillarum</i> in closed and open corrals through time at Cayo Diablo.	20
Table 3 The number of restocked <i>Diadema antillarum</i> in closed and open corrals through time at Los Lobos.....	20
Table 4 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition at the control stations between sites (Si), quadrat types (Qu: fixed or random), and sampling dates (Da).....	26

Index List

Index 1 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.	34
Index 2 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Los Lobos, Fajardo. .	34
Index 3 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in Ramicrusta spp. between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.	34
Index 4 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in Ramicrusta spp. between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.	35
Index 5 Photographs of permanent quadrats before and two months after the restocking of <i>Diadema antillarum</i> in Cayo Diablo and Los Lobos, Fajardo, Puerto Rico.	35
Index 6 Photographs of the grazing effects of <i>Diadema antillarum</i> two months after restocking in the corrals at Cayo Diablo and Los Lobos, Fajardo.....	39

Introduction

Over the past four decades, coral reefs in the Caribbean have dramatically changed, (Hughes 1994, Wilkinson 2008, Jackson et al. 2014). The abundance of reef-associated organisms, especially corals, has suffered a massive decline due to cumulative factors such as, hurricanes, disease outbreaks, bleaching, pollution, and overfishing (Bythell and Sheppard 1993, Bythell et al. 1993, Littler et al. 1993, Hughes 1994, Kramer et al. 2003). One of the most dramatic shifts in community structure occurred after the massive die-off of *Diadema antillarum*, a keystone herbivore. The 1983-1984 mass mortality of *D. antillarum* occurred throughout the Caribbean basin and was the most extensive and severe die-off ever recorded for a marine invertebrate (Lessios 1995). Before 1983, the presence of this organism was common (13-18 Ind m⁻²) on coral reefs in the in Puerto Rico (Bauer 1980, Vicente and Goenaga 1984), and played an important role in structuring coral reef communities by controlling algal abundance (Carpenter 1981, Carpenter 1986, Carpenter 1990a, Carpenter 1990b, de Ruyter van Steveninck and Bak 1986, Odgen et al. 1973, Robertson 1987, Sammarco 1982), productivity (Williams 1990) and is one of the principal agents of bioerosion on reefs (Lidz and Hallock 2000, Bak et al. 1984, Scoffin et al. 1980). After the massive die-off, populations were drastically reduced by 95-100% in many Caribbean locations (Lessios 1995), and at the same time, fleshy macroalgal cover increased between 100% and 250% (Phinney et al. 2001). The absence of *D. antillarum* did not only influence the benthic algal productivity of coral reef communities, but it also impinged on the settlement of coral recruits.

Presently, the recovery of *D. antillarum* has been slow and even absent at many locations in the Caribbean (Lessios 2016). In Puerto Rico, there has been a modest recovery in the population of *D. antillarum* (Mercado-Molina et al. 2014), nevertheless, densities are still far below pre-mass mortality numbers (Lessios 2016). It has been proposed that either larval mortality and/or post-

recruitment mortality processes could be the main factors regulating the adult population size of *D. antillarum* (Karlson and Levitan 1990). In Puerto Rico, in particular, the former has been discarded as it has been shown that upstream sources of “settlement-ready” larvae for *D. antillarum* are available (Williams et al. 2010); therefore larval supply and survival do not seem to be inhibiting the recovery of these populations. Consequently, recruitment-limited processes, such as post-settler and/or juvenile mortality may be regulating the population dynamics of *D. antillarum* in Puerto Rico.

It is assumed that, after a disturbance and under the right circumstances, an ecosystem will recover (Nystrom and Folke 2001); however many reefs in the world, and in particular in the Caribbean, have lost their capacity to recover from recurrent disturbances and have undergone long-term phase shifts (Hughes et al. 2003). The most spoken about phase-shift in scientific literature is the coral to fleshy macroalgal shift (Hughes 1994, Shulman and Robertson 1996, McClanahan and Muthiga 1998, Rogers and Miller 2006). Reefs characterized in permanent states of algal dominance usually signifies a loss of resiliency (Hughes et al. 2007) because macroalgal assemblages limit coral settlement, affect sediment deposition, and alter chemical properties close to the benthos (Birrell et al. 2008). In Puerto Rico, a more recent and potential threat to the coral reefs is the increasing abundance of a red encrusting algae, *Ramicrusta* sp. *Ramicrusta* is an encrusting red alga from the Peyssonneliaceae family (Rhodophyta). Currently there are three species of *Ramicrusta* identified in the Caribbean; *textilis*, *bonairensis*, and *monensis* (Ballentine et al. 2016). *Ramicrusta* forms a thin, crustose layer that spreads over the substrate and can grow over the living tissues of other organisms (Pueschel and Saunders 2009, Eckrich and Engel 2013, Ballentine et al. 2011, Ruiz 2015). In Bonaire, Eckrich and Engel 2013) observed the maximum overgrowth rates to range between 0.06 mm d⁻¹ and 0.08 mm d⁻¹. The rapid growth of *Ramicrusta*

makes this algae not only a threat to slow growing sessile-benthic organisms but has the potential to reduce the area of suitable substratum for coral settlement. Species of corals that can be negatively impacted by the expansion of *Ramicrusta* include the ESA listed species of *Acropora* and *Orbicella* (Williams pers comm, Ruiz 2015).

During 2016, *Ramicrusta* cover was high (41-75%) on Fajardo, Culebra Island, and Vieques reefs (Garcia Sais et al. 2016). By 2017-18 surveys, *Ramicrusta* was observed at 70% of the 42 monitoring sites. Moreover, *Ramicrusta* sp. was the main benthic substrate observed on shallow-water coral reefs at a majority of the reefs surveyed in Culebra Island and Fajardo. On the east coast, the abundance of *Ramicrusta* was astoundingly high and dominated the benthic composition at the majority of sites and depths, with the exception at the deeper sites at Culebra and Fajardo. Even more concerning was that *Ramicrusta* was more abundant than coral at 8 out of the 9 sites on the east coast and the cover was more than double that of coral cover at half of these sites. During 2018, I observed *Ramicrusta* overgrowing *Orbicella annularis* and *Acropora cervicornis*, among other corals at reefs in Culebra (see Figure 1).

Diadema antillarum, a keystone herbivore, has been witnessed eating *Ramicrusta* (Williams pers comm, Ruiz 2015, Williams 2017) and in the field I observed a noticeably lower cover of *Ramicrusta* in areas where *D. antillarum* were present. The goal of this project was to increase herbivory rates on two coral reefs in Fajardo and decrease algal abundance, specifically *Ramicrusta* sp. In addition, it is also expected, that coral recruitment abundance will increase as an indirect consequence of *D. antillarum* herbivory effect (Edmunds and Carpenter 2001).

Methods

This project was developed and implemented with the funds of National Oceanographic and Atmospheric Administration (NOAA) Restoration Center. They co-sponsored the project, as funds from DNER were not available until June 25, 2018.

Collection




Figure 1. Settlement sites in La Parguera, Puerto Rico

The settlement of *D. antillarum* is patchy and at times hard to identify and collect. We followed the methodology of Williams et al. (2010, 2011) settlement studies. They concluded that shelf-edge sites at La Parguera, Puerto Rico, Old Buoy and El Hoyo were “hot spots” for *D. antillarum* settlement (max of 1,064 Ind m⁻²). The supply of *D. antillarum*

settlors for this restoration experiment were collected at the Old Buoy (Fig. 1). The depth of the site ranges from 18-21 meters.

The reef substrate at Old Buoy is fairly flat with relatively low coral cover and diversity. Historically, there have been few adult *D. antillarum* recorded at both of these sites (<0.01 Ind m⁻²).




Figure 2 Astroturf or artificial turf used as settlement

Thirty mooring lines were placed at Old Buoy. On each mooring line, a rope containing 20, 8 × 8 cm pieces of artificial turf (Fig. 2) was attached to the cement blocks and mooring buoy (600 plates in total). The cement blocks were placed in the middle of a sand

channel, and not impacting the corals and reef on either side. The buoys were located at 6 m of depth and did not represent any navigational hazard. Plates were positioned in 9.8 meters of depth in the water column. During collections, the rope was detached and placed in a dry bag and a new rope with clean plates was attached. Settlement plates were replaced each month on twenty lines. On the other ten lines, plates were replaced every two months from June to September 2017 (total of five samplings). Plates were brought back to the laboratory to analyze.

Grow-out

Settlers (Fig. 3) were picked off each settlement plates, counted and transferred to a 45-gallon and 60-gallon tank. Both tanks were connected to a closed-filtering system. The closed system helped reduce sediment and other larvae from entering and settling in the tanks, while maintaining water quality (salinity and temperature). Tanks were cleaned once a week and one-




Figure 4 *Diadema antillarum* settler

third of the water was replaced with fresh seawater that was filtered and run through a UV light. Water quality measurements were recorded every week. These measurements included salinity,



Figure 3 Wet laboratory at the University of Puerto Rico, Mayaguez.

pH, temperature, nitrate, and ammonium. Settlers were transferred to wet tables or raceways (Fig. 4) once they reached a size of 5 mm in test diameter. Raceways were connected to a semi-closed circulating system. This allowed the water to be recycled through the tanks even if fresh seawater was not being

supplied. The system was flushed once a week and fresh saltwater entering the system is filtered through a sock. The sock was replaced and cleaned once a week and raceways were cleaned once every two weeks. Algae were collected and placed in each raceway every three to four days. Algal species such as, *Acanthomorpha*, *Chaetomorpha*, *Padina*, *Stypopodium*, and *Dictyota* spp., were collected in the field once a week and kept in a holding tank for *Diadema* feed. *D. antillarum* juveniles were transplanted to the reef once they reached a test size of at least 3-4 cm in test diameter.

Restocking

D. antillarum juveniles were transplanted to the backreef of Cayo Diablo (18.36038, -65.53325) and Los Lobos (18.37405, -65.57025) in Fajardo, Puerto Rico (Figure 5). This was the first attempt to restocked urchins on the east side of Puerto Rico. Six corrals were installed at each reef at 5-7 meters and held into place with rebar. The diameter of each corral was approximately 2m². Corrals were made of galvanized chicken wire with a 1-inch diameter mesh size. The plastic chicken wire was attached to the bottom of the corral to mold to the reef. Corrals were placed in the sand around isolated *O. annularis* colonies. At each reef, tops were installed on three of the corrals to enclose (hereafter closed) the area fully, and three corrals did not have tops. Given the site location and weather conditions on the east coast, monitoring did not occur as frequent as the restocking of *Diadema* in La Parguera during 2016. Therefore, fully enclosing the corrals permitted the escape of restocked *Diadema* and reassured the measurements of grazing effects. Monitoring took place one day, one week, one and two-month after the reintroduction. Individuals were counted inside and outside the corrals. Corrals were removed after the two month monitoring on October 23, 2018.




Figure 5 Map of the restoration sites, Cayo Diablo and Los Lobos in Fajardo, Puerto Rico.

Six 25 cm x 25 cm quadrats, with three random and three fixed were placed and photographed inside of each corral to monitor benthic cover through time. Fixed quadrats were placed in areas with high algal cover, especially *Ramicrusta*. Nails will be used to mark the position of fixed quadrats, allowing for the estimation of change. In addition, change in benthic composition was also monitored outside the corrals (control) with three random and fixed quadrats. Percentage cover of algae was discriminated to the lowest possible taxonomic level. The photographs were examined in the laboratory and the relative percentage cover of sessile organisms was visually estimated. Observations on percentage cover of other benthic organisms were also recorded. Photographs were not taken at Los Lobos after one month of restocking because of a camera malfunction.




Figure 6 Pictures of the different corrals in Cayo Diablo (top six pictures) and in Los Lobos (bottom six), in Fajardo, Puerto Rico. Corral number is arranged from left to right.

Statistics

Three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) tests (Anderson 2001) were performed at each site (Cayo Diablo and Los Lobos) to examine corral type (open or closed), quadrat type (fixed and random), and date. The sites were examined

separately because the dataset was unbalanced, for example, the benthic composition was not measured one month after restocking at Los Lobos. Benthic cover was analyzed visually and without the use of any software. Corral 6 was not analyzed at each of the sites because all *D. antillarum* escaped after one week of restocking. Each PERMANOVA procedure was based on Bray-Curtis similarity measures. SIMPER test were run to identify the contribution of benthic categories to the overall differences between years with the random and fixed quadrats and at the different sites. A Principal Component Analysis (PCA) plots were produced to visualize differences in benthic assemblages between years.

PERMANOVA analyses were also run to examine the differences in *Ramicrusta* cover between the sites, corral type and dates. Again, separate analyses were tested for random and fixed quadrats. For the analyses of *Ramicrusta* cover (univariate), the similarity matrix was based on Euclidean distances. Euclidean distance measures for univariate PERMANOVA analyses produce sums-of-squares estimates equivalent to parametric ANOVA (Anderson, 2001), and allow the same methodological framework to be used for all community attributes.

Results and Discussion

Collection

Twenty lines were cut and stolen during the August collection. The lines were cut again in September, but we arrived at the site and were able to retrieve them as they floated away. Therefore, only 200 plates were analyzed during August. The mooring lines were removed during our collection in September, and we did not proceed with October collection because of weather (hurricanes). The total amount of settlers collected from June to September are listed below in the table (Table 1)

Table 1. The total amount of settlers (#) collected at each sites from June to September 2017.

Month	Settlers
June	259
July	1355
August	439
September	1126
Total	3179

Grow-out

There were some modifications this year because Hurricane Irma hit the island on September 5, 2017 and Hurricane Maria on September 20, 2017. Electricity was lost after Irma for about a week but most of the island was without electricity, especially Isla Magueyes for many months (about three months) after Maria. The university ran the generators for only eight hours a day. Many sea urchins started perishing in the tanks, possibly due to oxygen constraints and the daily temperature variation, since the water was not flowing throughout the night. On October 12, 2017, I decided to move all sea urchins to cages and hung these cages under the dock. A total of 1,367 sea urchins were placed in two cages (43% survivorship). Cages were made out of PVC tubes and screen. It was challenging to keep potential predators (mantis shrimps, worms, and crabs) outside of cages. On January 4, 2018, 505 sea urchins were removed from the cages and placed back in the raceways outside (37% survivorship).

Restocking

Survivorship

On August 22, 2018, at 1:00 PM, 480 young adults (3-4cm) were transferred to Cayo Diablo (240 in total) and Los Lobos (240 in total). Reintroducing *D. antillarum* during late

afternoon hours allows for adaption to reef conditions overnight. Twenty-five *D. antillarum* were placed in each corral (~6 Ind m⁻²). The rest of the *D. antillarum* were released freely on the reef.




Figure 7 Pictures of the lab-reared *Diadema antillarum* restocked in the corral at Cayo Diablo (top pictures) and Los Lobos (bottom pictures) reefs in Fajardo, Puerto Rico.

Most of the *D. antillarum* escaped the corrals a day after the restocking. More sea urchins were retained in the corrals at Cayo Diablo than in Los Lobos. Seventeen urchins (11%) escaped at Cayo Diablo and 43 urchins (28.7%) escaped at Los Lobos (Table 2 and 3). Most of the escaped urchins were placed back into the fully enclosed corrals after one week. It was hard locating the sea urchins that we released because they were well hidden and may have moved from the area.

However, we were able to locate some of the freed or escaped individuals one month after restocking because their grazing was visually evident on the surrounding reef. The top was removed on one corral (Corral 1) at Los Lobos after one week, and most of the *D. antillarum* escaped in this corral. It is unknown why the sea urchins escape the enclosures. They may be looking for a better shelter or food source.

Table 2 The number of restocked *Diadema antillarum* in closed and open corrals through time at Cayo Diablo.

Corrals with tops				Corrals without tops			
Date	Corral 1	Corral 2	Corral 5	Date	Corral 3	Corral 4	Corral 6
8/22/2018	25	25	25	8/22/2018	25	25	25
8/23/2018	25	25	25	8/23/2018	23	22	23
9/1/2018	20	15	16	9/1/2018	4	4	0
9/22/2018	16	12	11	9/22/2018	2	3	0
10/23/2018	9	0	8	10/23/2018	3	1	1

Table 3 The number of restocked *Diadema antillarum* in closed and open corrals through time at Los Lobos.

Corrals with tops				Corrals without tops				
Date	Corral 1	Corral 4	Corral 5	Date	Corral 1	Corral 2	Corral 3	Corral 6
8/22/2018	25	25	25	8/22/2018		25	25	25
8/23/2018	21	25	25	8/23/2018		12	19	22
9/1/2018		12	12	9/1/2018	5	1	4	0
9/22/2018		10	12	9/22/2018	2	1	2	0
10/23/2018		8	11	10/23/2018	0	1	2	0

As seen in Tables 2 and 3, the presence of *D. antillarum* through time varied between the type of corrals (closed or open). Overall, the presence in corrals was more significant in the closed corrals than in the open corrals. After the first month, an average of 44% and 52% of the restocked individuals were recorded in fully closed corrals in Los Lobos and Cayo Diablo, respectively. While after the first month, only 5% and 6% were recorded in open corrals at Los Lobos and Cayo Diablo. I did notice that the condition of the corrals after the first month significantly deteriorated. There were a lot of holes, and some of the plastic nettings was torn off. Small (damselfish and bluehead wrasse) to medium (surgeonfish) fish were observed many times inside the closed

corrals. Two large spotted morays were counted in Corral 1 at Cayo Diablo and Corral 4 in Los Lobos by the second month of monitoring. It is unclear if the moray is a predator of *D. antillarum*. There were only two dead sea urchins observed during this study, which occurred in a fully enclosed Corral 1 at Cayo Diablo. The cause of death of these sea urchins is unknown.

The numbers of *D. antillarum* located in the corrals decreased even more so by the second month of monitoring. The majority of the corrals were destroyed and open (Fig. 8) by the second month. This was evident at Cayo Diablo and could be due to the wave or swell energy at this site since it is more exposed. No dead urchins, piles of spines, or skeletons were observed outside the corrals or the surrounding reef area during the entire monitoring. As witnessed by Williams et al. (2015), escaped individuals can move fast and far, about 20 to 30 meters from restocked areas. Therefore, it was theorized that the *D. antillarum* which could not be located inside the corrals were well hidden and/or far from the restocking area.



Figure 8 Photographs taken during the two-month monitoring showing the destructions of corrals at Cayo Diablo (top pictures) and Los Lobos (bottom pictures) reefs in Fajardo, Puerto Rico.

Benthic Cover

Pre-restocking

Before restocking, a visual assessment of *D. antillarum* populations was conducted at Cayo Diablo and Los Lobos. The density of *D. antillarum* around the corrals was low ($< 1 \text{ Ind m}^{-2}$), with the exception at Los Lobos, where high densities of *D. antillarum* adults ($\sim 4 \text{ Ind m}^{-2}$) were observed about 50 m north of the corrals at shallower depths ($< 2\text{m}$).




Figure 9 Photographs at the control station in Cayo Diablo (two photographs above random quadrat) and in Los Lobos (two photographs below fixed quadrats) in Fajardo, Puerto Rico.

The control station at Cayo Diablo, benthic algae was the dominant benthic substrate, with macroalgae (*Dictyota* spp.) and *Ramicrusta* contributing to the highest mean cover (mean \pm se) in the fixed ($56.7 \pm 9.4\%$) and random quadrats ($63.3 \pm 9.3\%$), respectively. At Los Lobos, *Ramicrusta* was the main contributor to the overall algal composition, with mean cover ranging from $39.7 \pm 10.2\%$ in random quadrats and $31.3 \pm 3.2\%$ in permanent quadrats.

In the corrals, the benthic composition was similar to that in the control station and was mainly characterized by the high cover of *Ramicrusta* at both Cayo Diablo and Los Lobos, ranging from $41.8 \pm 11.3\%$ to $54.2 \pm 11.0\%$ and from $62.8 \pm 9.3\%$ to $83 \pm 5.6\%$, respectively (Fig. 9). Turf algae were also prominent outside and inside corrals. Turf was both sparse and filamentous and or in thick mats (algal mats), which allowed for the capture of sand and other sediments. Inside the

corrals, the cover of other benthic organisms was low, especially that of coral cover, which ranged in cover between $0.11 \pm 0.1\%$ to $13.9 \pm 8\%$ at Cayo Diablo to 0% to $12.75 \pm 6.7\%$ at Los Lobos.




Figure 10 The mean cover of benthic organisms in the corral type (open and closed) and quadrat type (fixed and random) through the sampling events at Cayo Diablo, Fajardo.




Figure 11 The mean cover of benthic organisms in the corral type (open and closed) and quadrat type (fixed and random) through the sampling events at Los Lobos, Fajardo.



Restocking effects

The benthic composition did not vary at the control stations through the monitoring period (Table 4). Algal cover, specifically *Ramificrusta* and turf algae, continued to be the dominant substrate outside the corrals through the monitoring period. There was no grazing observed in the control area. Therefore, it can be assumed that the changes observed in the benthic cover inside the corrals, both open and closed, are due to the grazing of restocked *D. antillarum*.

Table 4 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition at the control stations between sites (Si), quadrat types (Qu: fixed or random), and sampling dates (Da).

Source	df	SS	MS	Pseudo-F	p value
Si	1	6414.4	6414.4	4.8184	0.015
Qu	1	2834.7	2834.7	2.1294	0.106
Da	3	3336.1	1112	0.83533	0.59
SixDa	2	1865.5	932.76	0.70068	0.618
SixQu	1	1614.9	1614.9	1.2131	0.331
QuxDa	3	3336.1	1112	0.83533	0.601
SixQuxDa	2	1865.5	932.76	0.70068	0.596

One week after the restocking, the benthic substrate was characterized by more turf algae, and less *Ramicrusta* (see Fig. 12). *Ramicrusta* was reduced by as much as 57% at Cayo Diablo and 43% at Los Lobos. Also, there was also a lower abundance of fleshy macroalgae, specifically *Dictyota* spp, after one week. Even though the abundance of *Ramicrusta* was reduced after one week (see Fig. 13), the reduction was not significant for all corals. Therefore, resulting in the insignificant statistical results of the overall grazing effects of *D. antillarum* on the benthic composition. These results are similar to the restocking efforts in 2016 at Media Luna in La Parguera, Puerto Rico (Williams 2017). Grazing was evident after one week, however only ~20% was effectively grazed of algae after one week. Differences in the benthic composition might have been more significant if there was higher retention of restocked individuals in the corals.






Figure 12 Photographs of the fixed quadrats before and one week after the restocking of *Diadema antillarum* at Cayo Diablo (first two sets of photographs) and Los Lobos (third set of photograph) in Fajardo, Puerto Rico.

Benthic change was more evident in the closed corrals, given there was more grazing pressure. The abundance of *D. antillarum* recorded in an open corral was low and the retention ranged from one to three individuals ($< 1 \text{ Ind m}^{-2}$). The grazing pressure was still evident in the open corrals, however it was limited to the area of homing, thus explaining the lack of significance in the PERMANOVAs. Densities of one individual per square meter is not uncommon on coral reefs around the Caribbean. The majority of reefs are characterized by even lower densities. As seen in this study, the densities of individuals, $< 1 \text{ Ind m}^{-2}$, did not significantly impact the benthic composition after two months. However, this might have changed with a more extended monitoring period. Future *D. antillarum* restoration projects should focus on restoring populations greater than one individual per square meter.




Figure 13 Photographs taken to highlight the grazing of restored *Diadema antillarum* one week after the restocking at Cayo Diablo and Los Lobos in Fajardo, Puerto Rico.

As seen in the PERMANOVA tables (Indices 1-2) and the Principal Coordinate Analysis (PCA, Figure 14 and 15), there was a shift in benthic community structure after one and two months after the restocking (9/22/2018, 10/23/2018). The difference in benthic assemblage after one month was due to the reduction of *Ramicrusta* cover. The abundance of *Ramicrusta* was significantly reduced over the monitoring period (Indices 3-4). This pattern was valid in both the fixed and random quadrats at each site. In the closed corrals and during the two-month monitoring, *Ramicrusta* cover ranged from 18.9% to 32.7% at Cayo Diablo and from 33.8% to 45.7% at Los Lobos. The reduction in *Ramicrusta* was slightly more evident at Cayo Diablo than at Los Lobos; where *Ramicrusta* was reduced as much as 63% at Cayo Diablo (Fig. 16) and 61% at Los Lobos (See Indices 5 and 6). After one and two months, the substrate inside the corrals was characterized

by more available space and thin filamentous turf algae for corals and other benthic organisms to settle and colonize. Coral settlers were not observed during the monitoring period, which is not surprising since monitoring lasted for two months. The evidence presented in this study supports that the action of restocking of *Diadema antillarum* may be useful mitigation tool to control *Ramircrusta* and other algal abundance on the coral reefs.




Figure 14 Principal Coordinate Analysis on the benthic composition of closed (left) and open (right) corals at Cayo Diablo during the sampling periods. Algae represents all macroalgae.




Figure 15 Principal Coordinate Analysis on the benthic composition of closed (left) and open (right) corals at Los Lobos during the sampling periods. Algae represents all macroalgae.




Figure 16 Photographs of a fixed quadrat in Corral 5 at Cayo Diablo before, one week and two months after the restocking of *Diadema antillarum*.

Conclusions and recommendations

- Only a few dead sea urchins were witnessed during the monitoring period. Therefore, it was assumed that the sea urchins escaped the corrals. Some of the escaped sea urchins were located one month after the restocking. Therefore, the survivorship of the restocked sea urchins might be high.
- Fully enclosing corrals are more effective in retaining the sea urchins for the monitoring period. The sea urchins escape open corrals and this usually occurs during the first couple weeks.
- It is necessary to fully enclose the corrals in order to effectively assess the change in benthic composition.
- As witnessed by Dr. Williams in laboratory and in this study, *Diadema antillarum* consume *Ramicrusta* spp. and other algae that can be a nuisance on a coral reef, *Dictyota* spp.
- *D. antillarum* are effective in reducing the *Ramicrusta* on a coral reef.
- Grazing impacts on algal cover are significantly evident after one month of restocking. However, this time scale might vary depending on the amounts of sea urchins in a given space.
- Restocked densities must be greater than 1 Ind m⁻² in order to observe significant benthic change on a coral reef.
- Given the results of this study, the restocking of *D. antillarum* might be an effective mitigation tool in the reductions of algal cover, especially *Ramicrusta*.

Acknowledgements

This project was possible with the funding of the National Atmospheric and Oceanic Administration (NA17NOS4820037) and the Department of Natural and Environmental Resources. A big thanks to Katie Flynn, Orlando Espinosa, Milton Carlo, Manuel Olmeda, Luis Rodriguez, Liajay Rivera, Catalina Morales, Francisco González, Jaaziel Garcia, and Nicolle Lebrón for their help in the field and the laboratory. In addition, thank you to the Department of Marine Science, University of Puerto Rico for the facilities and all the employees who helped out with the project.

Indices

Index 1 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.

Cayo Diablo					
Source	df	SS	MS	Pseudo-F	p value
Co	1	8597.5	8597.5	7.6289	0.002
Qu	1	1830.5	1830.5	1.6243	0.193
Da	3	13688	4562.7	4.0487	0.002
CoxQu	1	3468.1	3468.1	3.0774	0.038
CoxDa	3	2708.4	902.79	0.80109	0.577
QuxDA	3	1977	659.01	0.58477	0.748
CoxQuxDA	3	5079.8	1693.3	1.5025	0.164

Index 2 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in benthic composition between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Los Lobos, Fajardo.

Los Lobos					
Source	df	SS	MS	Pseudo-F	p value
Co	1	4750.6	4750.6	5.2569	0.01
Qu	1	132.27	132.27	0.14636	0.865
Da	2	23099	11550	12.78	0.001
CoxQu	1	362.62	362.62	0.40127	0.689
CoxDa	2	187.53	93.763	0.10376	0.949
QuxDA	2	1662.8	831.39	0.91999	0.454
CoxQuxDA	2	1666.4	833.21	0.92201	0.463


Index 3 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in Ramicrusta spp. between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.





Cayo Diablo					
Source	df	SS	MS	Pseudo-F	p value
Co	1	2492.7	2492.7	3.2456	0.077
Qu	1	278.77	278.77	0.36298	0.542
Da	3	7577.8	2525.9	3.2889	0.026
CoxQu	1	87.728	87.728	0.11423	0.727
CoxDa	3	378.61	126.2	0.16433	0.908
QuxDA	3	1838.6	612.87	0.79799	0.483
CoxQuxDA	3	1982.3	660.77	0.86037	0.446


Index 4 The results of a three-way distance Permutational Multivariate Analyses of Variance (PERMANOVA) to detect changes in *Ramicrusta* spp. between corral type (Co: open or closed), quadrat types (Qu: fixed or random), and sampling dates (Da) at Cayo Diablo, Fajardo.


Source	df	Los Lobos			Pseudo-F	p value
		SS	MS			
Co	1	3842.2	3842.2	4.6252	0.042	
Qu	1	255.67	255.67	0.30777	0.567	
Da	2	22300	11150	13.422	0.001	
CoxQu	1	330.04	330.04	0.3973	0.511	
CoxDa	2	1180	590.02	0.71026	0.479	
QuxDA	2	1848.3	924.13	1.1125	0.315	
CoxQuxDA	2	1016.8	508.39	0.61199	0.523	

Index 5 Photographs of permanent quadrats before and two months after the restocking of *Diadema antillarum* in Cayo Diablo and Los Lobos, Fajardo, Puerto Rico.













Index 6 Photographs of the grazing effects of Diadema antillarum two months after restocking in the corals at Cayo Diablo and Los Lobos, Fajardo. These are individual photographs taken of areas where the substrate has been grazed by Diadema antillarum.









Literature review

Abelson A (2006) Artificial reefs vs coral transplantation as restoration tools for mitigating coral reef deterioration: benefits, concerns, and proposed guidelines. *Bulletin of Marine Science* 78:151-159

Alvarez-Filip L, Carricart-Ganivet JP, Horta-Puga G, Iglesias-Prieto R (2013) Shifts in coral-assemblage composition do not ensure persistence of reef functionality. *Scientific Reports* 3:3486

Aronson RB, Precht WF (2001) White-band disease and the changing face of Caribbean coral reefs. *Developments in Hydrobiology* 159:25-38

Ballantine DL, Ruiz H, Lozada C, Norris JN (2016) The genus *Ramicrusta* (Peyssonneliales, Rhodophyta) in the Caribbean Sea, including *Ramicrusta bonairensis* sp. Nov. and *Ramicrusta monensis* sp. Nov. *Botanica Marina* 59:

Ballantine DL, Ruiz H (2013) Unique red algal reef formation in Puerto Rico. *Coral Reefs* 32:411

Ballantine DL, Athanasiadis A, Ruiz H (2011) Notes on the benthic marine algae of Puerto Rico. X Additions to the flora. *Botanica Marina* 54: 292-302

Birrell CL, McCook LJ, Willis BL, Diaz-Pulido GA (2008) Effects of benthic algae on the replenishment of corals and the implications for the resilience of coral reefs. *Oceanography and Marine Biology: An Annual Review* 43:25-63

Chiappone M, Swanson DW, Miller SL, Smith SG (2002) Large-scale surveys on the Florida Reef Tract indicate poor recovery of the long-spined sea urchin *Diadema antillarum*. *Coral Reefs* 21:155-159

Doherty PJ, Dufour V, Galzin R, Hixon MA, Meekan MG, Planes S (2004) High mortality during settlement is a population bottleneck for a tropical surgeonfish. *Ecology* 85:2422-2428

Eakin MC, Morgan JA, Heron SF, Smith TB, Liu G, Alvarez-Filip, et al. (2010) Caribbean corals in Crisis: Record thermal stress, bleaching and mortality in 2005. *PLoS ONE* 5(11):e13969

Eckrich CE, Engel S (2013) Coral overgrowth by an encrusting red alga (*Ramicrusta* sp.): a threat to Caribbean reefs? *Coral Reefs* 32: 81-84

Edmunds PJ, Carpenter RC (2001) Recovery of *Diadema antillarum* reduces macroalgal cover and increases abundance of juvenile corals on a Caribbean reef. *Proceedings of the National Academy of Science USA* 98:5067-5071

Edmunds PJ, Elahi R (2007) The demographic of a 15-year decline in cover of the Caribbean reef coral *Montastraea annularis*. *Ecological Monographs* 77:3-18

Forsman ZH, Page CA, Toonen RJ, Vaughan D (2015) Growing coral larger and faster: micro-colony-fusion as a strategy for accelerating coral cover. *PeerJ* DOI 10.7717/peerj.1313

Garcia-Sais R, Esteves R, Williams S, Sabater-Clavell J, Carlos M (2016) Monitoring of Coral Reef Communities from Natural Reserves in Puerto Rico: 2016. Final Report submitted to the Department of Natural and Environmental Resources (DNER), U. S. Coral Reef National Monitoring Program.

Garcia-Sais R, Esteves R, Williams S, Sabater-Clavell J, Carlos M (2015) Monitoring of Coral Reef Communities from Natural Reserves in Puerto Rico: 2015. Final Report submitted to the Department of Natural and Environmental Resources (DNER), U. S. Coral Reef National Monitoring Program.

Garcia-Sais R, Esteves R, Williams S, Sabater-Clavell J, Carlos M (2012) Monitoring of Coral Reef Communities from Natural Reserves in Puerto Rico: 2012. Final Report submitted to the Department of Natural and Environmental Resources (DNER), U. S. Coral Reef National Monitoring Program.

Garcia-Sais J, Appeldoorn R, Battista T, Bauer L, Bruckner A, et al. (2008) The state of coral reef ecosystems of Puerto Rico. In The state of coral reef ecosystems of the United States and Pacific freely associated states: 2008. National Oceanic and Atmosphere Administration. NOAA Technical Memorandum NOS NCCOS 73.

Green DH, Edmunds PJ, Carpenter RC (2008) Increasing relative abundance of *Porites astreoides* on Caribbean reefs mediated by an overall decline in coral cover. Marine Ecology Progress Series 359:1-10

Hughes TP (1994) Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. Science 265:1547-1551

Hughes TP, Rodrigues MJ, Bellwood DR, Ceccarelli D, Hoegh-Guldberg O, McCook L, Moltchanivskyj N, Pratchett MS, Steneck RS, Willis B (2007) Phase shifts, herbivory, and the resilience of coral reefs to climate change. Current Biology 17:1-6

Hughes TP, Tanner JE (2000) Recruitment failure, life histories, and long-term decline of Caribbean corals. Ecology 81:2250-2263

Jackson JBC (1997) Reefs since Columbus. Coral Reefs 16:S23-S32

Jackson JBC, Donovan M, Cramer K, Lam V (2014) Status and trends of Caribbean coral reefs: 1970-2012. Global Coral Reef Monitoring Network, IUCN

Luckhurst BE, Luckhurst K (1978) Analysis of the influence of substrate variables on coral reef fish communities. Marine Biology 49:317-323

McClanahan TR, Muthiga NA (1998) An ecological shift in a remote coral atoll of Belize over 25 years. Environmental Conservation 25:122-130

Mercado-Molina AE, Ruiz-Diaz CP, Sabat AM (2014) Survival, growth, and branch production of unattached fragments of the threatened hermatypic coral *Acropora cervicornis*. Journal of Experimental Marine Biology and Ecology 457:215-219

- Miller MW, Kramer KL, Williams SM, Johnston L, Szmant AM (2009) Assessment of current rates of *Diadema antillarum* larval settlement. *Coral Reefs*. 28(2): 511-515
- Mumby PJ (2016) Stratifying herbivore fisheries by habitat to avoid ecosystem overfishing of coral reefs. *Fish and Fisheries* 17:266-278
- Nyström M, Folke C (2001) Spatial resilience of coral reefs. *Ecosystems* 4:406-417
- Perry CT, Steneck RS, Murphy GN, Kench PS, Edinger EN, Smithers SG, Mumby PJ (2015) Regional-scale dominance of non-framework building corals on Caribbean reefs affects carbonate production and future reef growth. *Global Change Biology* DOI: 10.1111/gcb.12792
- Perry CT, Murphy GN, Kench PS, Smithers SG, Edinger EN, Steneck RS, Mumby PJ (2013) Caribbean-wide decline in carbonate production threatens coral reef growth. *Nature Communications* 4:1402
- Planes S (2002) Dynamics and diversity in a complex ecosystem. Ed Sale PF. In *Coral Reef Fishes*. Academic Press, San Diego, USA.
- Planes S, Lecaillon G (2001) Caging experiment to examine mortality during metamorphosis of coral reef fish larvae. *Coral Reefs* 20:211-218
- Pueschel CM, Saunders GW (2009) *Ramicrusta textilis* nov. (Peyssonneliaceae, Rhodophyta), an anatomically complex Caribbean alga that overgrows corals. *Phycologia* 48: 480-491
- Ramírez-Mella JT, García-Sais JR (2003) Offshore dispersal of Caribbean reef fish larvae: how far is it? *Bulletin of Marine Science* 72:997-1017
- Roff G, Mumby PJ (2009) Global disparity in the resilience of coral reefs. *Trends in Ecology and Evolution* 27:404-413
- Rogers CS, Miller J (2006) Permanent ‘phase shifts’ or reversible declines in coral cover? Lack of recovery of two coral reefs in St. John, US Virgin Islands. *Marine Ecology Progress Series* 306:103-114
- Rogers CS, McLain LN, Tobias CR (1991) Effects of Hurricane Hugo (1989) on a coral reef in St. John, USVI. *Marine Ecology Progress Series* 78:189-199
- Ruiz H (2015) The ecology of the red algae *Ramicrusta textilis*, its dynamic with corals and the evaluation of possible management strategies to minimize its threat to coral reefs around Puerto Rico. Report PR-DNER
- Shulman MJ, Robertson DR (1996) Changes in the coral reefs of San Blas, Caribbean Panama: 1983 to 1990. *Coral Reefs* 15:231-236
- Soto-Santiago FJ, Irizarry-Soto E (2013) The sea urchin *Diadema antillarum* (Echinodermata, Equinoidea), algal cover and juvenile coral densities in la Parguera, Puerto Rico. *Cuadernos de Investigación UNED* 5:

- Van Oppen MJH, Oliver JK, Putnam HM, Gates RD (2015) Building coral reef resilience through assisted evolution. *Proceedings of the National Academy of Sciences of the United States of America* 112:2307-2313Soto
- Vermeij MJA, Marhaver KL, Huijbers CM, Nagelkerken I, Simpson SD (2010) Coral larvae move toward reef sounds. *PLoS ONE* 5(5): e10660
- Weil E, Croquer A, Urreiztieta I (2009) Temporal variability and impact of coral diseases and bleaching in La Parguera, Puerto Rico from 2003-2007. *Caribbean Journal of Science* 45:221-246
- Williams SM (2017) A novel approach to the restoration of *Diadema antillarum* on coral reefs in the Caribbean. *Reef Encounters* 31: 48-50
- Williams SM, García-Sais J, Yoshioka PM (2011) Spatial variation of *Diadema antillarum* settlement in La Parguera, Puerto Rico. *Bulletin of Marine Science* 87(3): 531-540
- Williams SM, Yoshioka PM, García-Sais J (2010) Recruitment pattern of *Diadema antillarum* in La Parguera, Puerto Rico. *Coral Reefs* 29: 809-812